

Jean-Marie Dufour
 January 21, 2003

**TIME SERIES ANALYSIS
 EXERCISES
 STOCHASTIC PROCESSES 3**

1. Let $T \subseteq \mathbb{R}$. Which condition must satisfy a family of joint distribution functions $F(x_1, \dots, x_n; t_1, \dots, t_n)$ defined for all finite subsets $\{t_1, \dots, t_n\} \subseteq T$, where $n \geq 1$, in order for these distributions to be the distributions of a stochastic process? [See Brockwell and Davis (1991, Theorem 1.2.1).]
2. Show that any even and positive-definite function $\gamma : \mathbb{Z} \rightarrow \mathbb{R}$ is the autocovariance function of a Gaussian stationary process. [See Brockwell and Davis (1991, Theorem 1.5.1).]
3. Consider the function $\gamma : \mathbb{Z} \rightarrow \mathbb{R}$ defined by

$$\begin{aligned}\gamma(k) &= 1, \quad \text{if } k = 0 \\ &= \rho, \quad \text{if } |k| = 1 \\ &= 0, \quad \text{otherwise.}\end{aligned}$$

Show this function is an autocovariance function if and only if $|\rho| \leq 0.5$. [See Brockwell and Davis (1991, Example 1.5.1).]

4. Let $(Z_t : t \in \mathbb{Z})$ be *i.i.d.* $N[0, \sigma^2]$ random variables, and let a, b , and c be real constants. Determine which ones of the following processes are second-order stationary. For each stationary process, determine its mean and autocovariance function.
 - (a) $X_t = a + b Z_t + c Z_{t-1}$
 - (b) $X_t = a + b Z_0$
 - (c) $X_t = Z_1 \cos(ct) + Z_2 \sin(ct)$
 - (d) $X_t = Z_0 \cos(ct)$
 - (e) $X_t = Z_t \cos(ct) + Z_{t-1} \sin(ct)$
 - (f) $X_t = Z_t Z_{t-1}$
5. Let $(Y_t : t \in \mathbb{Z})$ be a second-order stationary process such that $E(Y_t) = 0, \forall t$, and let a and b be real constants.

- (a) If $X_t = a + bt + s_t + Y_t$, where s_t is a periodic function with period 12, show that the process

$$Z_t = (1 - B)(1 - B^{12})X_t$$

is second-order stationary.

- (b) If $X_t = (a + bt)s_t + Y_t$, where s_t is a periodic function with period 12, show that the process

$$Z_t = (1 - B^{12})(1 - B^{12})X_t$$

is second-order stationary.

6. Let $(X_t : t \in \mathbb{Z})$ and $(Y_t : t \in \mathbb{Z})$ be second-order stationary processes uncorrelated with each other, i.e. such that $Cov(X_s, Y_t) = 0, \forall s, t$.

- (a) Show that the process $X_t + Y_t$ is second-order stationary
 (b) Find the autocovariance function of $X_t + Y_t$.

7. Consider the process

$$S_t = \mu + S_{t-1} + u_t, \quad t = 1, 2, \dots$$

where $S_0 = 0$ and u_1, u_2, \dots are *i.i.d.* random variables with mean zero and variance σ^2 .

- (a) Determine the mean and covariance function of the process S_t . Is this process strictly stationary? second-order stationary?
 (b) Show that the process $Y_t = (1 - B)S_t, t = 1, 2, \dots$ is strictly stationary. Compute its mean and autocovariance function.

References

BROCKWELL, P. J., AND R. A. DAVIS (1991): *Time Series: Theory and Methods*. Springer-Verlag, New York, second edn.